



UNIVERSITI PUTRA MALAYSIA

**STRUCTURE ELUCIDATION AND BIOLOGICAL ACTIVITY OF
DITHIOCARBAZATE DERIVATIVES, THEIR
SCHIFF BASE LIGANDS AND METAL COMPLEXES**

KHOO TENG JIN

FS 2008 52

**Structure Elucidation and Biological Activity of Dithiocarbazate Derivatives, Their
Schiff Base Ligands and Metal Complexes**

By

KHOO TENG JIN

**Thesis Submitted to the School of Graduate Studies,
Universiti Putra Malaysia in Fulfillment of the Requirements for the Degree of
Doctor of Philosophy**

October 2008



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirements for the degree of Doctor of Philosophy

**STRUCTURE ELUCIDATION AND BIOLOGICAL ACTIVITY OF
DITHIOCARBAZATE DERIVATIVES, THEIR SCHIFF BASE LIGANDS AND
METAL COMPLEXES**

By

KHOO TENG JIN

October 2008

Chairman : Professor Dr. Karen Badri

Faculty : Science, Department of Chemistry

Dithiocarbazate Schiff base ligands and their complexes has been an active field of research concentrating on synthesis and characterization and their transition metal complexes. This work focuses on the synthesis and characterization of new dithiocarbazate ligands of *S*-3-picolyldithiocarbazate, *N*-Pyridin-3-ylmethylene-*N*-[5-(pyridine-3-ylmethylsulfanyl)-[1,3,4]thiadiazol-2-yl]-hydrazine, *S*-4-picolyldithiocarbazate and *N'*-Pyridin-4-ylmethylenehydrazinecarbodithioic acid pyridine-4-yl methylester. The Schiff bases were fully characterized using various physico-chemical and spectro-chemical analyses and its crystal structure elucidated. Work on these new Schiff base ligands has paved the way for further research into their complexes. A total of 56 complexes and 12 Schiff base ligands were synthesized each containing different functionalities which are, 4-aminoacetophenone, furyl and carboxyl and pyridine.

The synthesized and fully characterized Schiff bases and their complexes were then evaluated for their biological activities against antimicrobial and cytotoxicities studies. From the antimicrobial studies, Schiff bases have wider range of activity against the bacterial strains as compared to its complexes while complexes that comprises of d^{10} metals has the best potential as antimicrobial agent. In the evaluation of its cytotoxicities against selected cancer cell lines, copper(II), platinum(II) and palladium(II) metal complexes are strongly active with lowest CD_{50} values which are comparable to commercially available drugs. In conclusion, both the Schiff bases and their complexes can be further studied for its mechanism of action and has high potential to be developed into a much potent chemotherapeutic agents.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan Ijazah Doktor Falsafah

**ELUCIDASI STRUKTUR DAN AKTIVITI BIOLOGI LIGAN BASE SCHIFF
DAN LOGAM KOMPLEKS DERIVASI DITHIOCARBAZAT**

Oleh

KHOO TENG JIN

Oktober 2008

Pengerusi : Professor Dr. Karen Badri

Fakulti : Sains, Jabatan Kimia

Kajian terhadap dithiocarbazat base Schiff dan logam kompleks adalah bidang yang giat aktif dan tertumpu kepada sintesis dan penentuan struktur. Penyelidikan ini tertumpu kepada sintesis and penentuan struktur ligan dithiocarbazat yang baru yakni, *S*-3-picolildithiocarbazat, *N*-Pyridin-3-ylmethylene-*N*-[5-(pyridin-3-ylmethylsulfanyl)-[1,3,4]thiadiazol-2-yl]-hydrazin, *S*-4-picolildithiocarbazat dan *N*⁷-Pyridin-4-ylmethylenhydrazincarbodithioic acid pyridin-4-ylmethylester. Struktur base Schiff baru ini tentukan sepenuhnya menggunakan pelbagai teknik analisis kimia-fisikal dan spektrokimia. Penyelidikan ini telah berjaya membuka peluang kepada kajian yang lebih lanjut terhadap base Schiff dan logam kompleks baru ini. Sejumlah 56 kompleks dan 12 ligan base Schiff telah disintesis dimana ianya terdiri daripada pelbagai kumpulan berfungsi iaitu, 4-aminasetophinon, furil, carboxil dan piridin.

Ligan base Schiff dan kompleks logam yang telah ditentukan sepenuhnya telah dikaji untuk aktiviti biologinya terhadap antimikrob dan sitotoksik. Keputusan antimikrob

menunjukkan base Schiff mempunyai aktiviti yang lebih meluas terhadap jenis bakteria berbanding kompleks logamnya tetapi kompleks logam yang terdiri daripada logam transisi d^{10} lebih berpotensi sebagai agen antimikrob. Dalam kajian sitotoksik terhadap sel kanser yang dipilih, kompleks-kompleks logam yang terdiri dari kuprum(II), platinum(II) dan palladium(II) didapati amat active dengan nilai CD_{50} yang amat rendah dan boleh dibandingkan dengan nilai ubat antikanser komersial. Dengan ini, boleh disimpulkan bahawa ligan base Schiff dan logam kompleksnya boleh dilanjutkan kajian untuk menentukan mekanisma tindakan dan kedua-duanya mempunyai potensi yang tinggi untuk dibangunkan sebagai agen kemoterapi.

ACKNOWLEDGEMENTS

My utmost appreciation and gratitude goes to Professor Dr. Karen Badri for her most insightful guidance, endless support and her limitless patience in waiting for this thesis to be completed and most of all of what she has sacrificed which is privileged only to her sons. Without her, I would not be able to reach this far. My deepest thanks to Dr. Mohammed Ibrahim Mohammed Tahir for his friendship, support and wonderful suggestion, without him there would be no fun and motivation in research. To Professor Dr. M.T.H. Tarafder, the person who started it all and I never look back, my deepest thank you for the encouragement and ever professional guidance at the very beginning. I am also very grateful to Dr. Andrew Cowley and Dr. David Watkin for providing me with a comfortable working environment in their laboratory and their great advice. Also, my thanks to Professor Dr. Bohari M. Yamin and all my fellow labmates who has helped me along the way.

This work would not have any significant if not for the support and love from my parents, sisters and my wife. Thank you all for allowing me to come this far. This work is dedicated to you all.



For my grandparents, parents, sisters, wife and supervisors

Possibly, just possibly,

this is the chemistry of love

I certify that a Thesis Examination Committee has met on 13 October 2008 to conduct the final examination of Khoo Teng Jin on his thesis entitled "Structure Elucidation and Biological Activity of Dithiocarbazate Derivatives, Their Schiff Base Ligands and Metal Complexes" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.


Members of the Thesis Examination Committee were as follows:

Taufiq Yap Yun Hin, PhD
Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Halim Shaari, PhD
Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Tan Wee Tee, PhD
Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Hapipah Mohd Ali, PhD
Professor
Chemistry Department
Universiti Malaya
(External Examiner)


HASANAH MOHD GHAZALI, PhD
Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 30 December 2008

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Karen Ann Crouse, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Mohammed Ibrahim Mohammed Tahir, DPhil

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

M.T.H Tarafder, PhD


Professor
Faculty of Science
Rajshahi University
(Member)

Bohari M. Yamin, PhD

Professor
Faculty of Science and Technology
Universiti Kebangsaan Malaysia
(Member)

Manaf A. Ali, PhD

Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)




HASANA H. M. GHAZALI, PhD
Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 12 FEB 2009

DECLARATION

I hereby declare that this thesis is based on my original work except for any quotations or citations, which have been duly acknowledged. I also declare that this thesis or any part of it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.



KHOO TENG JIN

Date: 26/2/09

LIST OF TABLES AND SCHEME

- 4.1: Crystallographic Data and Structure Refinement Details of *N*-Pyridin-3-ylmethylene-*N*-[5-(pyridine-3-ylmethylsulfanyl)[1,3,4]thiadiazol-2-yl]-hydrazine
- 4.2: Selected Bond Length (Å) and Angles (°)
- 4.3: Crystallographic Data and Structure Refinement details of *N'*-Pyridin-4-ylmethylenehydrazinecarbodithioic acid pyridine-4-ylmethylester
- 4.4: Selected Bond Length (Å) and Angles (°)
- 5.1: Synthesized Schiff bases derived from different dithiocarbazate
- 5.1.1: Microanalytical Data for the Schiff bases and Complexes containing:
- 5.1: Solvation of Chloride Ion of the Complexes
- 5.1.2: Molar Conductance and Magnetic Data for the Complexes Containing
- 5.1.3.1: Crystallographic Data and Structure Refinement Details of [Ni(S2pPy2C)₂]
- 5.1.3.2: Selected Bond Length (Å) and Angles (°) of [Ni(S2pPy2C)₂]
- 5.1.3.3: Crystallographic Data and Structure Refinement Details of [Zn(S2pPy2C)₂]
- 5.1.3.4: Selected Bond Length (Å) and Angles (°) of [Zn(S2pPy2C)₂]
- 5.1.3.5: Crystallographic Data and Structure Refinement Details of [Cd₂(S4p2ACP)₂Cl₂]
- 5.1.3.6: Selected Bond Length (Å) and Angles (°) of [Cd₂(S4p2ACP)₂Cl₂]
- 5.1.3.7: Crystallographic Data and Structure Refinement Details of [Zn₂(S4p2ACP)₂Cl₂]
- 5.1.3.8: Selected Bond Length (Å) and Angles (°) of [Zn₂(S4p2ACP)₂Cl₂]
- 5.1.3.9: Crystallographic Data and Structure Refinement Details of [Ni(Sb4AP)₂]
- 5.1.3.10: Selected Bond Length (Å) and Angles (°) of [Ni(Sb4AP)₂]



- 5.2.1: Qualitative Antimicrobial Assay (100 mg cm^{-3})^a
- 5.2.2: Quantitative Antimicrobial Assay (MIC, $\mu\text{g ml}^{-3}$)^b
- 5.2.3: Cytotoxic Activities of the Schiff Bases and their Metal Complexes



LIST OF FIGURES

- 1.1: Dithiocarbazate acid derivatives
- 1.2: Possible isomers of dithiocarbazates
- 1.3: *thione-thiol* tautomerism
- 1.4: Complexation mode of dithiocarbazate derivatives with metal ions
- 1.5: The structure of S-alkyl- β -N-(6-methylpyrid-2-yl)-methylenedithiocarbazate
- 1.6: The structure of bis{S-methyl- β -N-(6-methylpyrid-2-yl)-methylene
- 1.7: Mono-ligated acetate Copper(II) complex [9]
- 1.8: Bis-ligated Copper(II) complex [9]
- 4.1: *thione-thiol* tautomerism of S-3-picolyldithiocarbazate in solution
- 4.2: Assigned carbon number for ^1H and ^{13}C NMR
- 4.4: ORTEP Diagram of N-Pyridin-3-ylmethylene-N-[5-(pyridine-3-ylmethylsulfanyl)-[1,3,4]thiadiazol-2-yl]-hydrazine with Atomic Numbering Scheme (at 50% Probability Displacement Ellipsoids).
- 4.5: Crystal packing viewed through *ab* and *bc* plane
- 4.6: Structure and assigned carbon number of S-4-picolyldithiocarbazate
- 4.7: Structure and assigned carbon number of N'-Pyridin-4-ylmethylene hydrazinecarbodithioic acid pyridine-4-ylmethylester
- 4.8: ORTEP Diagram of N'-Pyridin-4-ylmethylenehydrazinecarbodithioic acid pyridine-4-ylmethylester with Atomic Numbering Scheme (at 50% Probability Displacement Ellipsoids).
- 4.9: Crystal packing viewed through *a* axis
- 5.1.3.1: ORTEP Diagram of $[\text{Ni}(\text{S2pPy2C})_2]$ with 50% Probability Displacement Ellipsoids and Atomic Numbering Scheme.
- 5.1.3.2: Crystal packing of $[\text{Ni}(\text{S2pPy2C})_2]$ viewed along *b* axis



- 5.1.3.3: ORTEP Diagram of $[\text{Zn}(\text{S2pPy2C})_2]$ with 50% Probability Displacement Ellipsoids and Atomic Numbering Scheme.
- 5.1.3.4: Crystal packing of $[\text{Zn}(\text{S2pPy2C})_2]$ viewed through b axis
- 5.1.3.5: ORTEP Diagram of $[\text{Cd}_2(\text{S4p2ACP})_2\text{Cl}_2]$ with 50% Probability Displacement Ellipsoids and Atomic Numbering Scheme.
- 5.1.3.6: Crystal packing of $[\text{Cd}_2(\text{S4p2ACP})_2\text{Cl}_2]$ viewed through a axis
- 5.1.3.7: ORTEP Diagram of $[\text{Zn}_2(\text{S4p2ACP})_2\text{Cl}_2]$ with 50% Probability Displacement Ellipsoids and Atomic Numbering Scheme
- 5.1.3.8: Crystal packing of $[\text{Zn}_2(\text{S4p2ACP})_2\text{Cl}_2]$ viewed through b axis
- 5.1.3.9: ORTEP Diagram of $[\text{Ni}(\text{Sb4AP})_2]$ with 50% Probability Displacement Ellipsoids and Atomic Numbering Scheme



LIST OF APPENDICES

- A 4.1: FT-IR Spectrum of *S*-3-picolylthiocarbamate
- A 4.2: Mass Spectra of *S*-3-picolylthiocarbamate
- A 4.3: ^1H NMR of *S*-3-picolylthiocarbamate
- A 4.4: ^{13}C NMR of *S*-3-picolylthiocarbamate
- A 4.5: TGA curve of *S*-3-picolylthiocarbamate
- A 4.6: DSC curve of *S*-3-picolylthiocarbamate
- A 4.7: Atom coordinates and U_{iso} or U_{eq} and thermal parameters of *N*-Pyridin-3-ylmethylene-*N*'-[5-(pyridine-3-ylmethylsulfanyl)-[1,3,4]thiadiazol-2-yl]hydrazine
- A 4.8: FT-IR Spectrum of *S*-4-picolylthiocarbamate
- A 4.9: FT-IR Spectrum of *N*'-Pyridin-4-ylmethylenediazinecarbodithioic acid pyridine-4-ylmethylester
- A 4.10: Mass Spectra of *S*-4-picolylthiocarbamate
- A 4.11: Mass Spectra of *N*'-Pyridin-4-ylmethylenediazinecarbodithioic acid pyridine-4-ylmethylester
- A 4.12: ^1H NMR of *S*-4-picolylthiocarbamate
- A 4.13: ^{13}C NMR of *S*-4-picolylthiocarbamate
- A 4.14: ^1H NMR of *N*'-Pyridin-4-ylmethylenediazinecarbodithioic acid pyridine-4-ylmethylester
- A 4.15: ^{13}C NMR of *N*'-Pyridin-4-ylmethylenediazinecarbodithioic acid pyridine-4-ylmethylester
- A 4.16: DSC curve of *S*-4-picolylthiocarbamate
- A 4.17: DSC curve of *N*'-Pyridin-4-ylmethylenediazinecarbodithioic acid pyridine-4-ylmethylester
- A 4.18: TGA curve of the crude mixture before separation
- A 4.19: Atom coordinates and U_{iso} or U_{eq} and thermal parameters of *N*'-Pyridin-4-ylmethylenediazinecarbodithioic acid pyridine-4-ylmethylester



TABLE OF CONTENTS

	Page
ABSTRACT	ii
ABSTRAK	iv
ACKNOWLEDGEMENT	vi
APPROVAL	viii
DECLARATION	x
LIST OF TABLES	xi
LIST OF FIGURES	xiii
LIST OF APPENDICES	xv
CHAPTER	
1. INTRODUCTION	1
2. LITERATURE REVIEW	
2.1 Multidentate Nitrogen-Sulfur Donor Ligands their Metal Complexes and its Crystal Structures	6
2.2 Transition Metal Complexes Derived from Dithiocarbazate Schiff Base Ligands and their Biological Properties	12
2.3 Dithiocarbazate Schiff base Ligands and Their Peculiar Properties	15
3. MATERIALS AND METHODOLOGY	
3.1 Chemicals	19
3.2 Preparation of Ligands	20
3.3 Preparation of Schiff Bases and Metal Complexes	22
3.4 Physico-chemical Measurements and Elemental Analysis	23
3.5 Determination of Biological Activity	26
4. STUDIES OF NEW DITHIOCARBAZATE LIGANDS	
<i>S</i> -3-picolylthiocarbazate and <i>N</i> -Pyridin-3-ylmethylene- <i>N</i> -[5-(pyridine-3-ylmethylsulfanyl)-[1,3,4]thiadiazol-2-yl]-hydrazine	
4.1 Synthesis Methodology	28
4.2 Results and Discussion	
4.2.1 Characterization of <i>S</i> -3-picolylthiocarbazate by CHNS, FTIR, Mass Spec., ¹ H and ¹³ C NMR and thermal studies by TGA and DSC	29
4.2.2 Structure elucidation of <i>N</i> -Pyridin-3-ylmethylene- <i>N</i> -[5-(pyridine-3-ylmethylsulfanyl)-[1,3,4]thiadiazol-2-yl]-hydrazine by SXRD	32



<i>S</i> -4-picolyldithiocarbazate and <i>N</i> ⁷ -Pyridin-4-ylmethylene hydrazinecarbodithioic acid pyridine-4-ylmethylester	
4.3 Synthesis Methodology	35
4.4 Results and Discussion	
4.4.1 Characterization of <i>S</i> -4-picolyldithiocarbazate and <i>N</i> ⁷ -Pyridin-4-ylmethylene hydrazinecarbodithioic acid pyridine-4-ylmethylester by CHNS, FTIR, Mass Spec., ¹ H and ¹³ C NMR and thermal studies by DSC and TGA	36
4.4.2 Structure elucidation of <i>N</i> ⁷ -Pyridin-4-ylmethylene hydrazinecarbodithioic acid pyridine-4-ylmethylester by SXRD	40
4.5 Discussion and Conclusion	44
5. RESULTS AND DISCUSSIONS	
5.1 STUDIES OF DITHIOCARBAZATE SERIES CONTAINING DIFFERENT FUNCTIONALITIES	46
5.1.1 Microanalytical Data for the Schiff Bases and Their Complexes	48
5.1.2 Molar Conductance and Magnetic Data for the Complexes	54
5.1.3 X-ray Crystallographic Structure Elucidation	
5.1.3.1 The X-ray Crystal Structure of [Ni(S2pPy2C) ₂] and [Zn(S2pPy2C) ₂] complexes	58
5.1.3.2 The X-ray Crystal Structure of [Cd ₂ (S4p2ACP) ₂ Cl ₂] and [Zn ₂ (S4p2ACP) ₂ Cl ₂] complexes	66
5.1.3.3 The X-ray Crystal Structure of [Ni(Sb4AP) ₂] complex	75
5.2 BIOLOGICAL ACTIVITIES	
5.2.1 Qualitative and Quantitative Antimicrobial Activities of the Schiff Bases and Their Complexes	79
5.2.2 Cytotoxic Activities of the Schiff Bases and Their Complexes	89
6. CONCLUSION	94
REFERENCES	96
APPENDICES	103
PUBLICATIONS	116
BIODATA OF STUDENT	163
ACCOMPANYING CD	164

1. INTRODUCTION

Schiff bases are a diverse class of ligands containing O, N as well as S as their main donor atoms for chelation [1]. With this simple property, they have opened a wide and active field of studies in inorganic chemistry. This field of research continues to prosper because of continuous discovery of applications in a wide variety of fields specifically pharmaceuticals, agricultural chemicals and electronics. A more defined scope of research concentrates on synthesis and characterization of complexes containing nitrogen-sulfur donor ligands with transition metals and their properties. In many cases, there are significant differences in bioactivity after complexation.

Schiff bases of dithiocarbazate acid derivatives are one of these types of ligands. The structure is shown in Figure 1.1 where R = methyl, benzyl or pyridine.




Figure 1.1: Dithiocarbazate acid derivatives

In the solid state, dithiocarbazates were found to exist only as *cis-trans* and *trans-cis* isomers [2, 3] although theoretically they can evolve as four different types of isomers as shown in Figure 1.2. Conformation is assigned with respect to the N-C and C-S bond.




Figure 1.2: Possible isomers of dithiocarbazates

The types of isomers isolated are expected to be determined by the method of preparation and temperature of recrystallization. The *cis-trans* isomer was formed by recrystallization from benzene/chloroform solvent at room temperature while *trans-cis* was formed by recrystallization from methanol-water solvent in the ratio of 1:3 at -20°C . Using S-methyldithiocarbazate as an example, X-ray diffraction studies found the *cis-trans* conformers crystallize as a monoclinic, the space group $C 2/c$ with $Z' = 8$. The basic units of these structures are centrosymmetric dimers formed by $\text{NH}\cdots\text{N}$ hydrogen bonds. The component monomers are in the *cis-trans* conformation, where the group NH-NH_2 bends toward C-S and the SMe towards the opposite side. In a second conformer (*trans-cis*) of S-methyldithiocarbazate, the crystal has been obtained by crystallisation at low temperature and is triclinic with $Z' = 2$. The structure comprises centrosymmetric dimers linked by $\text{N-H}\cdots\text{S}$ hydrogen bonds, in contrast to the *cis-trans*

conformer where N-H·····N bonds are present. The vibrational spectra of both conformers differ markedly, but the change of conformation does not significantly alter the bond distances and angles of either structure. In solution, dithiocarbazate derivatives undergo thione-thiol tautomerism depending on the pH of the solution (Figure 1.3), while in the solid state, only the thione tautomer exists.




Figure 1.3: thione-thiol tautomerism

Dithiocarbazate derivatives have four donor atoms which are available to chelate with metal ions. Only two may ligate at a given time. Therefore, there are two possible complexation modes which are sulfur and β -nitrogen (NS) modes or both sulfur atoms (SS) mode (Figure 1.4). However, there are no examples of complexes of Schiff bases derived from *S*-alkyl ester of dithiocarbazic acid in which both sulfur atoms are coordinated to a metal ion.




Figure 1.4: Complexation modes of dithiocarbazate derivatives

A wide series of compounds with interesting chemical properties can be derived by substituting the –R group using R-halides while larger Schiff base molecules can be derived by using varieties of organic aldehydes or ketones in the condensation reaction. Interesting properties and geometries can be modified by chelating to different transition metals. These fascinating properties of the ligands forming different geometries, crystal morphologies and biological properties even though their molecular structures differ only slightly gave us the motivation for further research.

This thesis will discuss the synthesis of new ligand types containing 2-, 3- and 4-isomers of picoline substituted for –R, Schiff bases derived from these compounds through condensation reaction with different carbonyl compounds and complexation of the derivatives with transition metals Ni(II), Cu(II), Zn(II), Cd(II), Pd(II) and Pt(II). This will be followed by discussion of the characterization of the compounds and structure elucidation using various physico-chemical and spectroscopic techniques and the Single Crystal X-ray Diffraction (SXRD) methods when suitable single-crystals could be isolated. The thermal and electrochemical behaviors in solid and solution are also explored as well as their biological properties (qualitatively and quantitatively). Selected functional groups containing aminoacetophenone, fural, carbonyl, pyrrole and pyridine were introduced into the Schiff bases to study the effect of the presence of different functional groups in relation to biological activity.

It is hoped that this project will give a new dimension to this field of research and complement work done previously by our group and by other groups in addition to opening up new avenue for research in this area.

In more specific the objectives of this research are:

- (a) To synthesize new ligands derived from dithiocarbazate containing isomers of pyridine.
- (b) To synthesize new series of Schiff bases containing aminoacetophenone, fural, carbonyl, pyrrole and pyridine.
- (c) To synthesize complexes with transition metals of Ni(II), Cu(II), Zn(II) Cd(II), Pd(II) and Pt(II).
- (d) To characterize and elucidate the structure of the synthesized compound using various physico-chemical, spectroscopic and Single Crystal X-ray Crystallographic (SXRD) method.
- (e) To evaluate the qualitative and quantitative antimicrobial activities of the Schiff bases and their complexes.
- (f) To evaluate the cytotoxicity of the synthesized compound against cancer cell lines and
- (g) To correlate the biological activity with the structure of the synthesized compound.

2. LITERATURE REVIEW

2.1 Multidentate Nitrogen-Sulfur Donor Ligands their Metal Complexes and Crystal Structures

One of the least understood properties of this type of ligand is their chelating property which ranges from bidentate to multidentate. Although other physico-chemical methods are helpful in determining multidentate chelating capabilities, the best way to understand its chelating properties is by Single Crystal X-ray Diffraction (SXRD). The number of donor atoms in the Schiff base depends on the type of aldehyde or ketone condensed with the dithiocarbazic acid. A good representative of this type of ligation is in the bis{S-methyl- β -N-(6-methylpyrid-2-yl)-methylenedithiocarbazato}nickel(II) complex [4]. The uninegatively charged tridentate ligand formed by the condensation reaction between 6-methyl-2-formylpyridine with S-methyl and S-benzyl dithiocarbazates has the structure shown in Figure 1.5 [4].




Figure 1.5: The structure of S-alkyl- β -N-(6-methylpyrid-2-yl)-methylenedithiocarbazate

In the above case, the presence of 6-methylpyridine gives the ligand tridentate character through ligation of the mercaptide sulfur, azomethine nitrogen and the nitrogen

of the pyridine (NNS⁻) upon coordination with Ni(II) metal. The free Schiff bases does not display the $\nu(\text{SH})$ band at $\sim 2700 \text{ cm}^{-1}$ but exhibits $\nu(\text{NH})$ bands in the range $3100\text{--}3150 \text{ cm}^{-1}$ indicating that in the solid state it exists as a thioketo tautomer. Attempts by the author and his group to yield a mono-ligated Ni(II) complex with Cl⁻, Br⁻ or NCS⁻ as counter ion using large excess of Ni(II) salt always yielded the bis-ligated complex. This is in contrast to behavior seen for other related tridentate Schiff bases derived from 2-formylpyridine and S-methyl or S-benzyl-dithiocarbazate [5] which readily yielded the mono-ligated Ni(II) complex with a formula [Ni(NNS)X] where X = Cl, Br, NCS and I. The coordination modes of both complexes are determined by means of infrared spectra. Both types of complexes shows a shift in azomethine $\nu(\text{C}=\text{N})$ bands of the ligands to lower frequency and the increase in frequency of the hydrazinic N-N bond as a consequence of reduction between lone pairs of electrons on the hydrazine nitrogen atoms, supporting coordination *via* azomethine nitrogen atom. Coordination modes are further explained by means of structure elucidation using SXRD. The crystal structure of bis{S-methyl- β -N-(6-methylpyrid-2-yl)-methylenedithiocarbazato}nickel(II) is a six-coordinate environment with two Schiff base ligands coordinated to the nickel(II) ion as tridentate NNS chelating agents as shown in Figure 1.6. While the Schiff base coordinates in the iminothiolate form, a negative charge generated upon deprotonation is delocalized over the C-N-N-C bonds as indicated by intermediate bond distances of N-N = 1.389(9) Å, C-N = 1.26(1) Å and N-C = 1.31(1) Å, with the primary effect on the C-N bonds. By using SXRD, we are given the privilege to study the geometry of complexes, as in this case, the ligands are coordinated in the meridional configurations (with two azomethine nitrogen atoms *trans* to each other and the pyridine and sulfur atoms *cis*). This *mer* configuration has also been observed in nickel(II) complexes of